

UNIVERSIDADE DE LISBOA
FACULDADE DE CIÊNCIAS
DEPARTAMENTO DE BIOLOGIA ANIMAL



Combined effects of climate change and methylmercury exposure on the ecophysiology of juvenile meagre

Mestrado em Ecologia Marinha

Sofia dos Reis Alves e Dias Francisco

Dissertação orientada por:

Professor Doutor Rui Rosa
(Universidade de Lisboa, Faculdade de Ciências, Departamento de Biologia Animal)

Doutor António Marques
(IPMA – Instituto Português do Mar e da Atmosfera)

2015

UNIVERSIDADE DE LISBOA
FACULDADE DE CIÊNCIAS
DEPARTAMENTO DE BIOLOGIA ANIMAL



Combined effects of climate change and methylmercury exposure on the ecophysiology of juvenile meagre

Mestrado em Ecologia Marinha

Sofia dos Reis Alves e Dias Francisco

Dissertação orientada por:

Professor Doutor Rui Rosa
(Universidade de Lisboa, Faculdade de Ciências, Departamento de Biologia Animal)

Doutor António Marques
(IPMA – Instituto Português do Mar e da Atmosfera)

2015

This work was presented in the following scientific meetings

Francisco, S.; Pimentel, M. S.; Paula, J.R; Rosa, I.; Madeira, V.; Repolho, T.; Marques, A.; Rosa, R.

Combined effects of climate change and methylmercury exposure on marine fish ecophysiology.

23-27th March 2015

Third International Symposium on Effects of Climate Change on the World's Oceans
Santos City, Brazil

Francisco, S.; Maulvault, A.L.; Pimentel, M. S.; Lopes, A. R.; Paula, J.R.; Rosa, I; Madeira, V.; Repolho, T.; Marques, A.; Rosa R.

Impact of methylmercury exposure on marine fish ecophysiology under ocean warming and acidification.

3-7th May 2015

SETAC Europe 25th Annual Meeting Barcelona.
Barcelona, Catalonia, Spain

Acknowledgements

I owe my gratitude to all those people who have made this dissertation possible.

Firstly, I would like to express my sincere gratitude to Dr. Rui Rosa, for accepting me as his student and for believing in me, for all your supervision, support, enthusiasm, and immense knowledge that made this thesis possible.

I would like to thank to Dr. António Marques, for accepting to co-supervise this work and for all the help, support and encouragement provided.

I am thankful to Marta Pimentel for helping me through my first steps in this thesis, Ana Rita Lopes for all her teachings in the laboratory and support and José Paula for all the help and advices throughout this work. Their companionship, encouragement and support were very important for this dissertation. I also have to thank Tiago Repolho, Vanessa Madeira e Inês Rosa who also helped me during this work.

I would like to thank to all my colleagues from Laboratório Marítimo da Guia for the good moments, their sympathy and friendship, which made this work a better experience.

I also want to thank IPMA's team for receiving me so well, in particular, Ana Luísa Maulvault for the companionship and Ricardo for all the help during my work there.

I would like to thank to all of my dearest friends that helped me stay positive through this year, especially to Carolina, Inês, Vanessa and Steven, for listening, caring and always showing support.

I also want to thank to my family for all the support they provided me through my entire life, without them, I wouldn't be here.

Lastly, I have to thank to Gonçalo, without his love, encouragement and support, I would not have finished this thesis.

Resumo

Na época pré-industrial as concentrações de dióxido de carbono (CO_2) atmosférico eram aproximadamente 280 ppm e atualmente excedem os 390 ppm. É estimado que no ano 2100 as concentrações de CO_2 atmosférico cheguem aos 420 a 940 ppm. Este aumento tem provocado um aumento global da temperatura dos oceanos, que até 2100 é esperado aumentar entre 0,3 °C e 4,8 °C, especialmente nas zonas costeiras. Este aquecimento afetará a performance dos animais marinhos em diferentes estados do ciclo de vida através de mudanças na sua fisiologia.

Simultaneamente, os oceanos estão a absorver CO_2 , alterando o seu balanço químico, causando uma diminuição no pH dos oceanos e provocando assim, a acidificação dos oceanos. Desde a época pré-industrial, o pH da superfície do oceano diminuiu cerca de 0,1 unidades e é esperado que diminua entre 0,13 a 0,42 unidades até 2100. No entanto, estes valores poderão ser mais elevados em zonas costeiras. Esta acidificação irá afetar diversos organismos como bivalves e peixes. Uma das alterações que este aumento de CO_2 poderá envolver nos peixes é um dispêndio energético acrescido para acomodar as necessárias adaptações fisiológicas.

Além do aquecimento e acidificação dos oceanos, os organismos marinhos têm ainda que lidar com outro fator de *stress* existente: a contaminação. Os ecossistemas marinhos estão a ser contaminados com diversos compostos tóxicos, como os metais. Um destes metais é o mercúrio que é libertado para o ambiente através de fontes naturais ou antropogénicas. Depois da revolução industrial, as emissões de mercúrio antropogénico aumentaram drasticamente e as previsões apontam para a continuação deste aumento. Atualmente as emissões de mercúrio são maioritariamente de origem antropogénica e parte das mesmas são depositadas no oceano. A transformação de mercúrio em metilmercúrio ocorre através de bactérias existentes nos sedimentos. O metilmercúrio bioacumula em organismos aquáticos e biomagnifica ao atravessar os diferentes níveis tróficos, aumentando assim a sua toxicidade e chegando a níveis muito elevados nos organismos que ocupam o topo da teia trófica. Para além disso, pode causar efeitos adversos nos peixes, como por exemplo, alterar a performance reprodutiva, aumentar o *stress* oxidativo e alterar a atividade enzimática do sistema nervoso central.

Os organismos marinhos terão de lidar com os efeitos do aquecimento, acidificação e contaminação dos oceanos simultaneamente. No entanto, pouco se sabe sobre as interações entre estes fatores nos seres vivos. Alguns estudos mostram que os impactos das alterações climáticas nas espécies marinhas podem ser agravados pela

poluição e que as interações entre a temperatura e a poluição por metais influencia a tolerância fisiológica de animais aquáticos ectotérmicos. Além disso, a acidificação do oceano também pode intensificar a toxicidade dos metais.

Com o objetivo de avaliar as respostas biológicas de corvinas juvenis (*Argyrosomus regius*), em relação à exposição ao aquecimento dos oceanos (+ 4 °C) e acidificação ($\Delta\text{pH} = 0,5$ unidades) e ao MeHg ($8.02 \text{ mg kg}^{-1} \text{ dw}$), foi determinado o impacto combinado destes fatores sobre o índice de condição de Fulton, grau de encefalização, níveis de acetilcolinesterase, resposta ao choque térmico, peroxidação lipídica e atividade das enzimas antioxidantes. É importante referir que *A. regius* é um peixe resistente e pode facilmente adaptar-se às alterações ambientais, no entanto, esta espécie é encontrada em zonas costeiras e estuários, especialmente na fase juvenil, onde é expectável que as alterações climáticas irão ter maior impacto no futuro. Sendo a corvina um peixe de elevado valor comercial, é importante avaliar os efeitos futuros que estes fatores de *stress* terão sobre esta espécie.

Os resultados indicam que o aquecimento, a acidificação e a contaminação de mercúrio promovem diversas alterações fisiológicas. A bioacumulação de mercúrio foi afectada principalmente pelo aumento da temperatura, sendo o fígado o tecido que acumulou mais mercúrio. No entanto, não ocorreram diferenças na taxa de sobrevivência e na condição de Fulton. Relativamente à taxa de encefalização, apenas foram observadas alterações nos organismos sujeitos a aquecimento e hipercapnia, onde o tronco cerebral aumentou sob a condição de aquecimento e pH atual.

Foi observado um aumento em ambas as condições controlo de temperatura (18°C) e pH (8.0) na resposta das proteínas de choque térmico. No entanto, ocorreu uma diminuição significativa destas proteínas, sob condições de aquecimento e hipercapnia. Em oposição, os níveis de malondialdeído observados não mostraram diferenças significativas entre os tratamentos. Relativamente às enzimas de *stress* oxidativo, a atividade da glutathione S-transferase (GST) não se alterou com a contaminação, mas o aquecimento provocou um aumento na atividade de GST. Sob contaminação, a GST apresentou níveis mais elevados nos organismos provenientes do tratamento controlo (temperatura e pH) quando comparado com os organismos do tratamento de aquecimento e hipercapnia. A atividade da catalase não apresentou diferenças significativas nos organismos dos tratamentos não contaminados, mas com a contaminação a enzima aumentou nos peixes sujeitos à temperatura controlo e hipercapnia. Por último, a atividade da superóxido dismutase (SOD) não apresentou diferenças nos peixes dos tratamentos não contaminados, mas apresentando níveis mais elevados nos organismos sujeitos a contaminação no tratamento de temperatura controlo e hipercapnia.

Este estudo parece indicar que a corvina conseguirá adaptar-se às condições climáticas expectáveis no futuro em ambientes contaminados com MeHg. O aquecimento será o fator que mais afetará a acumulação de mercúrio na corvina e algumas variáveis irão ser afetadas pela contaminação. Embora a corvina tenha sofrido algumas alterações a nível fisiológico, estas não foram nocivas para o peixe. Em conclusão, estes resultados indicam, que a corvina poderá adaptar-se às futuras condições climáticas mesmo sob contaminação de mercúrio.

Palavras-chave:

Corvina, alterações climáticas, mercúrio, bioacumulação, encefalização, neurotransmissão colinérgica, *stress* oxidativo.

Abstract

The increase in anthropogenic carbon dioxide (CO₂) in the atmosphere is leading to higher ocean temperature and lower pH. At the same time, mercury released into the environment is also a fact, being deposited in the ocean. Thus, marine animals are being challenged by ocean warming, acidification and contamination simultaneously. However, their biological and physiological responses to these environmental changes still remain unknown. This study evaluated meagre (*Argyrosomus regius*) responses to the effects of ocean warming (+4°C), acidification ($\Delta\text{pH} = 0.5$ units) and MeHg exposure (8.02 mg kg⁻¹ dw), on Fulton condition index, degree of encephalization, acetylcholinesterase (AChE) levels, heat shock response (HSR), lipid peroxidation [malondialdehyde (MDA) concentration] and antioxidant enzyme activities [glutathione S-transferase (GST), catalase and superoxide dismutase (SOD)]. Temperature was the factor leading to an increase of MeHg accumulation in meagre. Survival and Fulton condition were not affected by any of the factors. Encephalization coefficient was only affected under the warming and hypercapnia non-contaminated treatment and only brain stem increased in response to warming. Moreover, AChE was only influenced under contaminated treatments. HSR only varied in muscle tissue and was mostly affected by warming treatments (non-contaminated and contaminated). In contrast, MDA levels were not influenced by warming, hypercapnia or contamination. GST increased under the warming non-contaminated treatment. Catalase was higher in samples from the contaminated hypercapnia treatment compared to the corresponding non-contaminated treatment, whereas SOD reached higher levels under the hypercapnia contaminated treatment. The present study shows that meagre will not suffer major consequences under predicted future ocean contaminated conditions. Although meagre suffered physiological alterations under these conditions, these responses were not harmful for the fish. These results suggest that meagre will be able to adapt to the predicted future ocean conditions even under mercury contamination. However, future studies will be necessary to comprehensively understand how marine biota will respond to the combined effects of future warming, acidification and pollution.

Keywords:

Meagre, climate change, mercury, bioaccumulation, encephalization, cholinergic neurotransmission, oxidative stress

Table of contents

Acknowledgements	iii
Resumo	iv
Palavras-chave:	vi
Abstract	vii
Keywords:	vii
List of figures and tables	x
1. Introduction	1
1.1. Global warming	1
1.2. Ocean acidification	2
1.3. Mercury contamination	3
1.4. Synergisms	4
1.5. Stress biomarkers.....	5
2. Objectives	7
3. Material and methods	8
3.1. Collection of specimen and incubation.....	8
3.2. Diet.....	9
3.3. Total mercury and methylmercury accumulation.....	10
3.4. Survival and Fulton condition.....	11
3.5. Brain sampling	11
3.6. Acetylcholinesterase (AChE) activity.....	12
3.7. Heat shock response, lipid peroxidation and antioxidant enzyme activities....	12
3.7.1. Preparation of tissue extracts:.....	12
3.7.2. Heat shock proteins.....	13
3.7.3. Lipid peroxides assay (malondialdehyde concentration).....	13
3.7.4. Glutathione S-Transferase (GST) activity	14
3.7.5. Catalase (CAT) activity	15
3.7.6. Superoxide dismutase (SOD) activity.....	15
3.8. Statistical analyses.....	16
4. Results	17
4.1. Total mercury accumulation.....	17
4.2. Survival, Fulton condition and encephalization coefficient.....	19

4.3.	Macro areas coefficient.....	20
4.4.	Brain macro areas acetylcholinesterase activity.....	22
4.5.	Heat shock response and lipid peroxidation.....	25
4.6.	Antioxidant enzymes activities.....	27
5.	Discussion	29
5.1.	Total mercury accumulation.....	29
5.2.	Fulton condition and survival	30
5.3.	Encephalization.....	30
5.4.	Brain macro areas	31
5.5.	Acetylcholinesterase (AChE) activity.....	32
5.6.	Oxidative stress	33
6.	Final remarks.....	36
	References	37
	Supplemental material	48

List of figures and tables

Figure 1. Components of the climate system that would be expected to change in a warming world exhibit trends consistent with warming (arrow direction denotes the sign of the change) (IPCC, 2013).....	1
Figure 2. Linkages between the buildup of atmospheric CO ₂ and the slowing of coral calcification due to ocean acidification (Hoegh-Guldberg et al., 2007).....	2
Figure 3. Temperature effects on the thermal windows of aerobic performance on aquatic animals. Synergistic stressors like ocean acidification (by CO ₂) narrow thermal windows (Pörtner and Farrell, 2008).	3
Figure 4. Global budgets for preindustrial and current mercury cycling in oceans (Sunderland and Mason, 2007).....	4
Figure 5. Meagre (<i>Argyrosomus regius</i>).	8
Figure 6. Fish brain macro areas: telencephalon, optic tectum, cerebellum, diencephalon and brainstem (Gonçalves et al., 2008)	11
Figure 7. Impact of ocean acidification (Δ pH 0.5) and warming (+4 °C) on total mercury accumulation on A) brain B) gills, C) muscle and D) liver of juvenile meagre (<i>Argyrosomus regius</i>) exposed to methylmercury for 30 days. Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. Capital letters represent significant differences between tissues in the same treatment.	18
Figure 8. Impact of ocean acidification (Δ pH 0.5), warming (+4 °C) and methylmercury contamination on A) Fulton condition index and B) the encephalization coefficient of juvenile meagre (<i>Argyrosomus regius</i>). Values represent mean+s.d. Different letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition.	19
Figure 9. Impact of ocean acidification (Δ pH 0.5), warming (+4 °C) and methylmercury contamination on macroareas coefficient between the five areas of the brain A) telencephalon, B) optic tectum, C) diencephalon, D) cerebellum and E) brain stem of juvenile meagre (<i>Argyrosomus regius</i>). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition.	

Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. 21

Figure 10. Impact of ocean acidification ($\Delta\text{pH } 0.5$), warming ($+4\text{ }^{\circ}\text{C}$) and methylmercury contamination on the acetylcholinesterase activity on the five areas of the brain A) telencephalon, B) optic tectum, C) diencephalon, D) cerebellum and E) brain stem of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. Capital letters represent significant differences between tissues in the same treatment..... 24

Figure 11. Impact of ocean acidification ($\Delta\text{pH } 0.5$), warming ($+4\text{ }^{\circ}\text{C}$) and methylmercury contamination on HSP in A) gills, B) muscle and C) liver and on D) muscle MDA concentration in juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Capital letters represent significant differences in HSP between tissues in the same treatment..... 26

Figure 12. Impact of ocean acidification ($\Delta\text{pH } 0.5$), warming ($+4\text{ }^{\circ}\text{C}$) and methylmercury contamination in the A) GST activity, B) Catalase activity and C) SOD activity in the muscle tissue of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Different small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. 28

Table 1. Composition of the two diets (i.e. dry inert pellets) used for juvenile meagre feeding..... 9

Table 2. Total mercury and methylmercury concentrations in the non-contaminated and contaminates feed 10

Table 3. Laboratory performance on standard reference materials for total mercury and methylmercury (mg kg^{-1} dry weight)..... 11

1. Introduction

1.1. Global warming

In the preindustrial era the atmospheric carbon dioxide (CO₂) concentrations were approximately 280 ppm (Caldeira and Wickett, 2003; IPCC, 2013) and are now exceeding 390 ppm (IPCC, 2013). By the year 2100, it is expected that atmospheric CO₂ concentrations reach values between 420 and 940 ppm, depending on the emissions scenario (IPCC, 2014). This increase in CO₂ (and other greenhouse gases) is causing a rise in temperature and therefore global warming (IPCC, 2013) (Fig. 1). Researchers predict a temperature rise between 0.3 °C to 4.8 °C by the year 2100 (IPCC, 2013). Sea surface temperature (SST) has significantly warmed during the past 30 years and it is predicted that will continue to increase, especially in coastlines (IPCC, 2014). Warming affects the performance of marine organisms at different stages in their life history through changes in physiology, morphology and behavior (Harley et al., 2006). Species with ample thermal windows, short generation times and a range of genotypes in populations will be favored under warming conditions (Pörtner and Farrell, 2008). Ultimately, species thermal tolerance and adaptability will determine the distribution and abundance of species under these conditions (Harley et al., 2006).




Figure 1. Components of the climate system that would be expected to change in a warming world exhibit trends consistent with warming (arrow direction denotes the sign of the change) (IPCC, 2013).

1.2 Ocean acidification

Simultaneously, oceans are absorbing CO_2 , changing their chemical balance (IPCC, 2013). Dissolved CO_2 increases the bicarbonate ion (HCO_3^-) and causes a decrease in carbonate ion (CO_3^{2-}) (IPCC, 2013). Ocean CO_2 uptake causes a decrease in ocean pH, resulting in ocean acidification (IPCC, 2013). Since the preindustrial time, ocean surface water pH has decreased 0.1 units and it is expected to decrease up to 0.13-0.42 units until 2100 (IPCC, 2014), however these values may be higher in coastal areas (IPCC, 2014). Ocean acidification will have impacts especially on calcifying organisms due to the reduction in seawater pH and carbonate ion concentrations which causes calcium carbonate (CaCO_3) saturation (Azevedo et al., 2015; Hoegh-Guldberg et al., 2007; Orr et al., 2005) (Fig 2). The reduction of carbonate-ions concentrations diminishes the rate of calcification of marine organisms weakening carbonate skeletal structures (Raven et al., 2005). Besides calcifying species, ocean acidification will also have an impact in other organisms, such as fish (Ishimatsu et al., 2008). High $p\text{CO}_2$ may involve an extra energy expenditure for physiological adaptations, especially in acid-base regulation (Ishimatsu et al., 2008). It has also been demonstrated that hypercapnia causes alterations in fish biological responses like aerobic scope (Munday et al., 2009; Rummer et al., 2013), decreases growth and thermal tolerance (Checkley et al., 2009; Pimentel et al., 2014b) and behavioral changes (Cripps et al., 2011; Munday et al., 2010; Pimentel et al., 2014a).




Figure 2. Linkages between the buildup of atmospheric CO_2 and the slowing of coral calcification due to ocean acidification (Hoegh-Guldberg et al., 2007).

These global ocean changes are expected to cause negative responses in marine organisms, essentially because of the synergistic effects between these two factors.

These effects have already been reported by various studies in diverse marine biota. Investigators demonstrated that in marine organisms, the synergism between warming and acidification has a higher negative effect on calcification, survival, and reproduction than when stressors act in isolation (Harvey et al., 2013). Predicted ocean warming and acidification caused shorter embryonic periods, lower survival rates and enhanced premature hatching in the cuttlefish and have drastic effects in survival rates of summer embryos of a coastal squid (Rosa et al., 2014b; Rosa et al., 2013). Pimentel et al., (2015) demonstrated that the effects on several physiological and biochemical variables in flatfish larvae change greatly under both ocean warming and acidification. Munday et al., (2009) also reported that aerobic scope and mortality in reef fish are affected by ocean warming and acidification. Pörtner and Farrell, (2008) presented a conceptual model of how synergistic stressors like ocean acidification (by CO₂) narrow thermal windows of aerobic performance according to species-specific sensitivities (Fig. 3).




Figure 3. Temperature effects on the thermal windows of aerobic performance on aquatic animals. Synergistic stressors like ocean acidification (by CO₂) narrow thermal windows (Pörtner and Farrell, 2008).

1.3 Mercury contamination

In addition to global warming and ocean acidification, marine biota is also dealing with another stress factor: pollution. Marine ecosystems are becoming contaminated with several toxic compounds including metals. One of these metals is mercury, which is a very toxic element, as well as its compounds (Korbas et al., 2011). Mercury releases into the environment can occur through natural and anthropogenic sources (e.g. coal combustion, industrial activities, mining) (Nriagu and Pacyna, 1988; Richter et al., 2014). After the industrial revolution, mercury anthropogenic emissions increased drastically (Sunderland and Mason, 2007) and predictions say it is likely they will continue to increase (Streets et al., 2009). Nowadays mercury emissions are mostly of

anthropogenic origin and part of these emissions are deposited in the ocean sediment (Selin et al., 2008; Sunderland and Mason, 2007) (Fig. 4).




Figure 4. Global budgets for preindustrial and current mercury cycling in oceans (Sunderland and Mason, 2007).

Bacteria present in aquatic sediments transform the inorganic mercury form into methylmercury (MeHg), which is highly toxic (Richter et al., 2014). Due to the biochemical conditions that lead the Hg transformation in MeHg by bacteria, estuaries and coastal areas are favorable zones for this transformation to occur (Dijkstra et al., 2013). This compound bioaccumulates in aquatic organisms and goes through biomagnification in aquatic food webs, increasing its toxicity and reaching elevated concentrations in high trophic level biota (Campbell et al., 2005; Evers et al., 2011). Hence fish consumers, including humans, are at high risk of exposure to MeHg (Richter et al., 2014). MeHg comprises around 90–95% of the total mercury (HgT) in fish (Burger et al., 2003; Gray et al., 2000). It can cause several adverse effects in these animals, for example, disturb reproductive performance (Hammerschmidt et al., 2002; Matta et al., 2001), enhance oxidative stress (Berntssen et al., 2003; Gonzalez et al., 2005; Keyvanshokoo et al., 2009; Larose et al., 2008; Mieiro et al., 2010), alter activity of enzymes of the central nervous systems (De Domenico et al., 2013; Gill et al., 1990) and change swimming behavior and predator avoidance (Weis and Weis, 1995; Zhou and Weis, 1998).

1.4 Synergisms

Although marine organisms will have to deal with the effects of ocean warming, acidification and contamination simultaneously, little is known about the interactions between climate change and contaminants on marine organisms. The impacts of climate

change on marine species can be aggravated by pollution (Noyes et al., 2009). Interactions between temperature and metal pollution influence the physiological tolerance to both stress factors in aquatic ectotherms (Sokolova and Lannig, 2008). Methylmercury bioaccumulation will increase at lower trophic levels under the predicted temperature rise, which consequently will increase MeHg levels throughout the food chain (Dijkstra et al., 2013). Elevated temperature exacerbates biological responses in freshwater fish exposed to metal contamination (Dorts et al., 2014; Lapointe et al., 2011; Sappal et al., 2014). Moreover, ocean acidification can also intensify metal toxicity (Han et al., 2014). For instance, it has been shown that acidification enhances the toxicity of copper in the polychaete *Arenicola marina*, showing that there are higher toxicity responses under reduced seawater pH (Campbell et al., 2014). Furthermore, temperature and pH affect the bioaccumulation of different metals in cephalopods early life stages (Lacoue-Labarthe et al., 2012; Lacoue-Labarthe et al., 2009; Lacoue-Labarthe et al., 2011).

Only few studies were conducted regarding climate change and contamination responses of marine organisms and, to my knowledge, none uses mercury as the metal contaminant. Most of them just evaluate the bioaccumulation in the organisms or visible responses such as mortality. Also, in fish, this type of response is usually analyzed in freshwater species. That said, there is a lack of knowledge in the response of marine biota, particularly in fish, to ocean warming, acidification and contamination simultaneously.

1.5 Stress biomarkers

As previously said, climate change can cause differences in the accumulation of mercury in fish (Dijkstra et al., 2013) plus, mercury has a great effect in the nervous system (Baatrup, 1991). In teleost fish, brain grows continuously during its lifetime (Ekström, 1994; Kotschal et al., 1998) and can be affected by environmental conditions such as habitat complexity or different rearing conditions especially in early stages (Kihlslinger and Nevitt, 2006; Pollen et al., 2007). Acetylcholinesterase (AChE) is an important enzyme in the nervous system because it hydrolyses the neurotransmitter acetylcholine (Romani et al., 2003). It has been reported that near-future carbon dioxide can alter neurotransmitter function in fish (Nilsson et al., 2012) and it is known that this enzyme is affected by mercury (Gill et al., 1990; Richetti et al., 2011). Nonetheless, no information exists on how ocean warming, acidification and mercury contamination will influence AChE response in fish.

Heat shock proteins (HSP) production occurs in response to high temperatures (Repolho et al., 2014; Rosa et al., 2014a; Rosa et al., 2012) and metal contamination

(Rajeshkumar and Munuswamy, 2011; Williams et al., 1996). High temperatures cause formation of oxygen reactive species (ROS) (Abele et al., 2002; Lesser, 2006) and HSP production helps repairing, refolding and eliminating damaged or denatured proteins and protect and control ROS formation (Sokolova et al., 2011). The increase in ROS may also cause lipid peroxidation, one of the most common cellular injury mechanisms (Lesser, 2006). ROS production induce antioxidant enzymes activity, such as superoxide dismutase (SOD), which converts superoxide (O_2^-) into hydrogen peroxide (H_2O_2); catalase (CAT) which converts H_2O_2 into water (H_2O) and oxygen (O_2) and glutathione S-transferase (GST), which is involved in the protection against xenobiotics (Lesser, 2006; Wang et al., 2000).

2. Objectives

The aim of this study was to evaluate the biological responses of juvenile meagre (*Argyrosomus regius*), to the effects of ocean warming (+4°C), acidification ($\Delta\text{pH} = 0.5$ units) and MeHg exposure ($8.02 \text{ mg kg}^{-1} \text{ dw}$). More specifically, I determined the combined impact of such climate-related variables and contamination on Fulton condition index, degree of encephalization, acetylcholinesterase levels, heat shock response, lipid peroxidation and antioxidant enzyme activities. It is worth noting that *A. regius* is a resilient fish and can easily adapt to changes in the environment (Monfort, 2010). However, this species is found in coastlines and estuaries, especially in the juvenile stage, where climate change and mercury will have more impact in the future. Meagre is a high commercial value fish (El-Shebly et al., 2007), thus being important to evaluate the future effects of these stressors on this species.

3. Material and methods

3.1. Collection of specimen and incubation

Juvenile organisms of *Argyrosomus regius* ($n = 85$; Fig. 5) (mean \pm standard deviation; total weight: 4.26 ± 2.8 g; total length: 6.30 ± 1.2 cm), from IPMA's aquaculture pilot station (EPPO, Olhão, Portugal) were transported to the aquaculture facilities of Laboratório Marítimo da Guia (LMG, MARE, Faculty of Sciences) in August 2014. The organisms were randomly placed in eight 90L tanks with separate recirculating aquaculture systems (RAS) filled with filtered natural seawater and equipped with glass wool, bio-balls and protein skimmers to maintain water quality.



Figure 5. Meagre (*Argyrosomus regius*).

After an acclimation period of 15 days, fish were subjected to ocean warming ($+4$ °C) and acidification ($\Delta\text{pH} = 0.5$ units) and MeHg contamination conditions, more specifically, the trial was divided in eight treatments: i) 19 °C, pH 8.0 (control conditions) and control feed (MeHg: $0.06 \text{ mg kg}^{-1} \text{ dw}$; HgT: $0.07 \text{ mg kg}^{-1} \text{ dw}$), ii) 19 °C, pH 8.0 and contaminated feed (MeHg: $8.02 \text{ mg kg}^{-1} \text{ dw}$; HgT: $8.28 \text{ mg kg}^{-1} \text{ dw}$), iii) 19 °C, pH 7.5 (control temperature and hypercapnic scenario) and control feed (MeHg: $0.06 \text{ mg kg}^{-1} \text{ dw}$; HgT: $0.07 \text{ mg kg}^{-1} \text{ dw}$), iv) 19 °C, pH 7.5 and contaminated feed (MeHg: $8.02 \text{ mg kg}^{-1} \text{ dw}$; HgT: $8.28 \text{ mg kg}^{-1} \text{ dw}$); v) 23 °C, pH 8.0 (warming scenario with control pH) and control feed (MeHg: $0.06 \text{ mg kg}^{-1} \text{ dw}$; HgT: $0.07 \text{ mg kg}^{-1} \text{ dw}$); vi) 23 °C, pH 8.0 and contaminated feed (MeHg: $8.02 \text{ mg kg}^{-1} \text{ dw}$; HgT: $8.28 \text{ mg kg}^{-1} \text{ dw}$); vii) 23 °C, pH 7.5 (warming and hypercapnic scenario) and control feed (MeHg: $0.06 \text{ mg kg}^{-1} \text{ dw}$; HgT: $0.07 \text{ mg kg}^{-1} \text{ dw}$); and viii) 23 °C, pH 7.5 and contaminated feed (MeHg: $8.02 \text{ mg kg}^{-1} \text{ dw}$; HgT: $8.28 \text{ mg kg}^{-1} \text{ dw}$), with ten to twelve fish per treatment, during 30 days. The MeHg exposure occurred through the feed. Fish were fed two to three times a day and the

quantity of food provided was approximately 1% of animal weight per day. Salinity was kept at 35.0 ± 1.0 g/L and the photoperiod was fixed to 12 h light: 12 h dark. Temperature in the tanks was controlled and maintained using chillers, and pH was maintained with a Profilux system. Temperature, pH and salinity were measured every day. Ammonia and nitrite were regularly monitored and kept within recommended levels. Seawater carbonate system speciation (Table S1) was calculated once a week from total alkalinity (determined according to Sarazin et al., (1999) and pH measurements). Total dissolved inorganic carbon (CT), $p\text{CO}_2$ and aragonite saturation were calculated using the CO2SYS software (Lewis et al., 1998), with dissociation constants from Mehrbach et al., (1973) as refitted by Dickson and Millero, (1987).

3.2. Diet

The non-contaminated and contaminated fish were fed with different diets. The composition of the two diets (Table 1) was similar diverging in one aspect: the contaminated diet was fortified with MeHg (inserted in the form of MeHg(II) chloride, CH_3ClHg , 99.8 %, Sigma-Aldrich, previously solubilized in ethanol). The pellet given to the fish in the contaminated treatment had approximately 8.02 mg kg^{-1} dw of MeHg and 8.28 mg kg^{-1} dw of HgT (Table 2).

Table 1. Composition of the two diets (i.e. dry inert pellets) used for juvenile meagre feeding.

Ingredients	%
Fishmeal LT70	28
Fishmeal 60	20
Fish soluble concentrate	2.5
Soy protein concentrate	5
Wheat gluten	5.5
Maize gluten	5
Soybean meal 48	9
Wheat meal	5.5
Whole peas	5
Fish oil	13.5
Vitamin and mineral premix	1
Dry matter (DM), %	94.3
Crude protein, %DM	51.7

Crude fat, %DM	18.9
Ash, %DM	9.6

Table 2. Total mercury and methylmercury concentrations in the non-contaminated and contaminates feed

Feed	HgT (mg kg ⁻¹)	MeHg (mg kg ⁻¹)
Non-contaminated	0.07 ± 0.00	0.06 ± 0.00
Contaminated	8.28 ± 0.10	8.02 ± 0.10

3.3. Total mercury and methylmercury accumulation

Methylmercury was extracted from the samples (fish and feeds) as described by Scerbo and Barghigiani, (1998), i.e. freeze-dried samples (approximately, 200 mg) were hydrolyzed in 10 ml of hydrobromic acid (47 % w/w, Merck), followed by MeHg extraction with 35 mL toluene (99.8 % w/w, Merck) and toluene removal with 6 ml cysteine aqueous solution (1 % L-cysteinium chloride in 12.5 % anhydrous sodium sulfate and 0.775 % sodium acetate; Merck). Then, total Hg and MeHg were determined in all samples (10-15 mg for solids or 100-200 µL for liquids) by atomic absorption spectrometry (AAS), following the method 7473 of the EPA, (2007), using an automatic Hg analyser (AMA 254, LECO, USA). Mercury concentrations were calculated from linear calibration (using, at least, five different standard concentrations), with a Hg(II) nitrate standard solution (1000 mg L⁻¹, Merck) dissolved in nitric acid (0.5 mol L⁻¹, Merck), and the detection limit was 0.005 mg kg⁻¹, wet weight (ww). Accuracy was checked through the analysis of the certified reference material DORM-4, and results obtained in the present study were within the certified range of values (Table 3). A minimum of three measurements (replicates) were performed per sample. Blanks were always tested in the same conditions as the samples. Prior to utilization, all laboratory ware was cleaned with nitric acid (20 % v/v) for 24h and rinsed with ultrapure water to avoid contamination. All standards and reagents were of analytical (pro analysi) or superior grade.

Table 3. Laboratory performance on standard reference materials for total mercury and methylmercury (mg kg⁻¹ dry weight).

	Standard reference material	HgT	MeHg
Present work	DORM-4*	0.390 ± 0.025	0.354 ± 0.031
Certified value		0.410 ± 0.055	0.353 ± 0.062

*fish protein certified reference material for trace metals, National Research Council Canada, Canada

3.4. Survival and Fulton condition

Survival rate was calculated as the percentage of live fish the end of the experiment, with respect to the number of fish at the beginning. After being euthanized fish were weighed and measured (TL). Fulton's condition (measure of individual fish's health) was calculated using the formula $K = (\text{weight}/\text{TL}^3) \times 100$.

3.5. Brain sampling

Fish were euthanized with an overdose of tricaine mesylate solution (MS222, Pharmaq; 500–1000 mg/L) and the spinal cord sectioned to be sure that the animal was dead. The brain was then visually macro-dissected from the cranial cavity, under a stereoscope. Afterwards, brain was dissected into five macro areas: telencephalon, diencephalon, optic tectum, cerebellum and brain stem (Fig. 6) and each area was weighted in an analytical Sartorius balance.




Figure 6. Fish brain macro areas: telencephalon, optic tectum, cerebellum, diencephalon and brainstem (Gonçalves et al., 2008)

3.6. Acetylcholinesterase (AChE) activity

Meagre brain areas were homogenized on ice in 1.0 mL of PBS solution (0.14M NaCl, 253 0.003M KCl, 0.01M Na₂HPO₄, 0.002M KH₂PO₄, pH 7.4) using a teflon tissue grinder for microtubes. The crude homogenates were then centrifuged (10 min at 14,000 x g at 4 °C). Then, 100 µl of supernatant were diluted to 1:2 in sodium phosphate buffer (50 mM, pH 8). The determination of acetylcholinesterase activity (EC 3.1.1.7; AChE) was performed based on an optimized Ellman method adapted to 96-well microplate (Ellman et al., 1961; Magnotti et al., 1987), in which thiocholine produced by the action of acetylcholinesterase forms a yellow (nitrobenzoate) color with 5,5'-dithiobis(2-nitrobenzoic acid), proportional to the AChE activity. All chemicals used in the assay were purchased to Sigma-Aldrich (Steinheim, Germany). Briefly, 50 µL of diluted samples, were added to each 96 microplate well (Greiner, Germany). Additionally, 50 µL of assay buffer were added to two microplate wells for blanks. Then an acetylthiocholine reaction mixture was prepared freshly in ultra-pure water [50 mM phosphate buffer; 75 mM Acetylthiocholine iodide solution; 1.0 mM of 5,5'-dithiobis(2-nitrobenzoic acid)] and 250 µL were added to each microplate well. Afterwards, the microplate was gently shaken for 30s at room temperature and absorbance read at 415 nm in a microplate reader (Bio-Rad, Benchmark model, USA) at every minute during 10 minutes. Then, AChE activity was calculated considering that one unit of enzyme catalyzes the production of 1.0 µmole of thiocholine per minute, under the assay conditions, and expressed as nmol min⁻¹ mg⁻¹ of total protein following normalization.

3.7. Heat shock response, lipid peroxidation and antioxidant enzyme activities

3.7.1. Preparation of tissue extracts:

Muscle, liver and gills samples (n = 6 per treatment) were homogenized (Ultra-Turrax, Ika, Staufen, Germany) in accordance to body mass of each sample in homogenization buffer, 300 mg tissue per 1 ml phosphate buffered saline solution (PBS, pH 7.3): 0.14 M NaCl, 2.7 mM KCl, 8.1 mM Na₂HPO₄, 1.47 mM KH₂PO₄] by using a glass/PTFE Potter Elvehjem tissue grinder (Kartell, Turin, Italy). All homogenates were then centrifuged (20 min at 14 000 g at 4 °C) and the antioxidant enzyme activities, lipid peroxidation and heat shock response (HSR) were quantified in the supernatant fraction. All enzyme assays were tested with commercial enzymes obtained from Sigma-Aldrich (St. Louis, MO, USA), and each sample was run in triplicate. The enzyme results were

normalize by measuring the total protein content of the samples according to the Bradford method (Bradford, 1976).

3.7.2. Heat shock proteins

Heat shock protein (HSP70/HSC70) content was assessed by Enzyme-Linked Immunoabsorbent Assay (ELISA) adapting a protocol from Njemini et al., (2005). Briefly, 10 μl of the homogenate's supernatant was diluted in 250 μl of PBS (1x), and 50 μl of the diluted sample was added to a 96-well microplates (Nunc brand, Roskilde, Denmark) and allowed to incubate overnight at 4 °C. In the next day, the microplates were washed (3X) in 0.05% PBS-Tween-20. One hundred microliters of blocking solution [1% bovine serum albumin (BSA) Sigma-Aldrich] was added to each well and left to incubate at room temperature for 2 h. After washing the 96-well plates, 50 μl of 5 $\mu\text{g ml}^{-1}$ primary antibody (anti-HSP70/HSC70, Acris, San Diego, CA, USA), detecting 72 and 73 kDa proteins corresponding to the molecular mass of inducible and constitutive isoforms, respectively, was added to each well and then incubated overnight at 4 °C. According to the manufacturer details the primary antibody HSP70/HSC70 (AM12032PU-N) has a broad range reactivity including diverse fish species. On the next day, the nonlinked antibody was removed by washing the microplates again, which were then incubated for 2 h at room temperature with 50 μl of 1 $\mu\text{g ml}^{-1}$ of the secondary antibody, antimouse IgG, Fab specific, alkaline phosphatase conjugate (Sigma-Aldrich). After three additional washes, 100 μl of substrate (SIGMA FASTTM p-Nitrophenyl Phosphate Tablets, Sigma-Aldrich) was added to each well and incubated 10-30 min at room temperature. Stop solution (50 μl ; 3 N NaOH) was added to each well, and the absorbance was then read at 405 nm in a 96-well microplate reader (Asys UVM 340, Biochrom, USA). The amount of HSP70/HSC70 present in the samples was calculated from a curve of absorbance based on serial dilutions of purified HSP70 active protein (Acris) to give a range from 0 to 2000 ng ml^{-1} . The results were expressed in relation to the sample total protein (ng mg^{-1} total protein).

3.7.3. Lipid peroxides assay (malondialdehyde concentration)

Lipid peroxides assay was determined by the quantification of a specific end-product of the oxidative degradation process of lipids, malondialdehyde (MDA). The thiobarbituric acid reactive substances (TBARS) protocol (Uchiyama and Mihara, 1978) was used. Briefly, 5 μl of each sample, treated as described before, were added to 45 μl of 50 mM monobasic sodium phosphate buffer, followed by adding 12.5 μl of sodium dodecyl sulfate (8.1%), 93.5 μl of trichloroacetic acid (20%, pH = 3.5) and 93.5 μl of

thiobarbituric acid (1%) to each microtube. Then, 50.5 µl of ultrapure water was added to this mixture and placed in a vortex for 30 s. Afterward, a needle was used to puncture the lids and microtubes were incubated in boiling water (10 min) followed by cooling on ice. Subsequently, 62.5 µl of ultrapure water and 312.5 µl of *n*-butanol pyridine (15 : 1, v/v) (Sigma-Aldrich, Hamburg, Germany) were added and microtubes centrifuged (5000 x g; 5 min.). Triplicates of 150 µl of the supernatant of each reaction were put into a 96-well microplate and absorbance was read at 532 nm. To quantify the lipid peroxides (MDA content), an eight-point calibration curve (0–0.3 µM TBARS) was calculated using malondialdehyde (dimethylacetal) (MDA) (Merck, Switzerland) standards. Results were expressed in relation to the sample total protein (nmol mg⁻¹ total protein).

3.7.4. Glutathione S-Transferase (GST) activity

Glutathione S-transferase (GST) total activity (EC 2.5.1.18) was determined according to the procedure described by Habig et al., (1974) and optimized for 96-well microplate (Sigma Technical Bulletin, CS0410). This assay uses 1-Chloro-2,4-dinitrobenzene (CDNB) as substrate and, upon conjugation of the thiol group of glutathione to the CDNB substrate, there is an increase in the absorbance. Therefore, the enzyme activity was determined spectrophotometrically by measuring the formation of the conjugates of glutathione (GSH) and 1-chloro-2,4-dinitrobenzene (CDNB). The assay contained 200 mM L-glutathione (reduced), Dulbecco's PBS and 100 mM 1-chloro-2,4-dinitrobenzene (CDNB) solution. Equine liver GST (Sigma-Aldrich) was used as positive control to validate the assay. Then, to perform the assay, 180 µl of substrate solution were added to 20 µl of GST standard or sample in each well of a 96-well microplate (Nunc-Roskilde) and the absorbance at 340 nm was recorded every minute for 6 min, using a plate reader (Asys UVM 340, Biochrom, USA). Thereby the increase in absorbance is directly proportional to GST activity. Then, the GST activity was calculated using a molar extinction coefficient for CDNB of 5.3 εmM (Sigma Technical Bulletin, CS0410), as follows:

$$GST\ activity = \frac{\Delta A_{340}/min}{0.0053} \times \frac{Total\ volume}{Sample\ volume} \times dilution\ factor$$

The result are expressed in relation to total protein of the sample (nmol min⁻¹ mg⁻¹ total protein)

3.7.5. Catalase (CAT) activity

Catalase activity was assessed through and adaptation of the method described by Johansson and Borg, (1988). In this assay, 20 μ l of each sample, 100 μ l of 100 mM potassium phosphate and 30 μ l of methanol were added to a 96-well microplate, which was promptly shaken and incubated for 20 minutes. Afterwards, 30 μ l of potassium hydroxide (10 M KOH) and 30 μ l of purpald (34.2 mM in 0.5 M HCl) were added to each well, and the plate shaken and incubated for another 10 minutes. Subsequently, 10 μ l of potassium periodate (65.2 mM in 0.5 M KOH) was added to each well and a final incubation was performed for 5 minutes. Using a microplate reader (BIO-RAD, Benchmark, USA), enzymatic activity was determined spectrophotometrically at 540 nm. Formaldehyde concentration of the samples was calculated based on a calibration curve (from 0 to 75 μ M formaldehyde), followed by the calculation of the CAT activity of each sample, where one unit of catalase is defined as the amount that will cause the formation of 1.0 nmol of formaldehyde per minute at 25 °C. The results are expressed in relation to total protein content (nmol min⁻¹ mg⁻¹protein).

3.7.6. Superoxide dismutase (SOD) activity

The SOD assay follows the nitro blue tetrazolium (NBT) method adapted from Sun et al., (1988). In this method, superoxide radicals (O_2^-) are generated by the reaction of xanthine with xanthine-oxidase (XOD), and reduce NBT to formazan. SOD competes with NBT for the dismutation of O_2^- into peroxide (H_2O_2) and molecular oxygen. The assay was performed using a 96-well microplate (Nunc-Roskilde), adding to each well 200 μ l of 50 mM phosphate buffer (pH 8.0) (Sigma-Aldrich), 10 μ l of 3 mM EDTA (Riedel-de Haën, Seelze, Germany), 10 μ l of 3 mM xanthine (Sigma-Aldrich), 10 μ l of 0.75 mM NBT (Sigma-Aldrich) and 10 μ l of SOD standard or sample. The reaction started with the addition of 100 mU XOD (Sigma-Aldrich) and the absorbance at 560 nm was recorded every minute for 5 minutes, using a plate reader (Asys UVM 340, Biochrom, USA) at 25 °C. A negative control included all components except SOD or sample, producing a maximal increase in absorbance at 560 nm, which allowed determining the inhibition percentage per minute, caused by SOD activity. SOD from bovine erythrocytes (Sigma-Aldrich) was used as standard and positive control. The SOD inhibition percentage was expressed in % inhibition mg⁻¹ of total protein.

3.8. Statistical analyses

Three-way ANOVAs (repeated measurements; using pH and temperatures and contamination) were conducted to detect significant differences in mercury accumulation, Fulton condition, acetylcholinesterase activity, heat shock response, lipid peroxidation and antioxidant enzyme activities. The differences in encephalization coefficient and macro areas coefficient were investigated in the following manner: for the encephalization coefficient a linear regression was made with the brain masses values of non-contaminated control conditions (19 °C pH 8.0), with these linear regression, predicted brain mass was calculated for the respectively fish mass of each fish of each condition. For the macro area coefficient five linear regressions were made with each brain macro area of non-contaminated control conditions (19 °C pH 8.0) and with these linear regressions, predicted macro areas masses were calculated for the respectively total brain mass of each fish and condition. Residuals were calculated with predicted mass value minus the observed mass and outliers were excluded using the ROUT method. Then, three-way ANOVAs (with temperature, acidification and contamination as variables) were conducted to detect significant differences in all variables measured. All statistical analyses were performed for a significance level of 0.05, using Statistica 12 software (StatSoft Inc., Tulsa, OK, USA).

4. Results

4.1. Total mercury accumulation

At the end of the 30-days trial, the warming and control pH condition elicited the highest mercury accumulation in most tissues analyzed (Fig. 7; $P < 0.05$). Under control conditions, gills and liver revealed higher levels of accumulation (6.25 mg kg^{-1} and 7.06 mg kg^{-1} , respectively) than muscle (3.74 mg kg^{-1}) ($P < 0.05$). In contrast, under the control temperature and hypercapnia treatment (19°C , pH 7.5), muscle accumulated less mercury (4.57 mg kg^{-1}) than liver (9.84 mg kg^{-1}) and the same happened under the warming and control pH condition ($P < 0.05$). Under the warming and hypercapnia scenario (23°C , pH 7.5) gills accumulated less mercury (3.25 mg kg^{-1}) than brain (5.29 mg kg^{-1}) ($P < 0.05$).




Figure 7. Impact of ocean acidification (Δ pH 0.5) and warming (+4 °C) on total mercury accumulation on A) brain B) gills, C) muscle and D) liver of juvenile meagre (*Argyrosomus regius*) exposed to methylmercury for 30 days. Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. Capital letters represent significant differences between tissues in the same treatment.

4.2. Survival, Fulton condition and encephalization coefficient

During the trial, no mortalities were registered in all treatments. Furthermore, Fulton condition (K) did not show any significant differences between treatments (Fig. 8A; $P > 0.05$). Warming and hypercapnia changed significantly the encephalization coefficient under the non-contaminated condition (Fig. 8B; $P < 0.05$). Under this condition, brain was significantly bigger than its predicted value. Under contaminated conditions, encephalization coefficient did not displayed significant differences ($P > 0.05$), however, it was significantly different between the non-contaminated and contaminated warming and hypercapnia scenarios ($P < 0.05$).





Figure 8. Impact of ocean acidification ($\Delta\text{pH } 0.5$), warming ($+4\text{ }^{\circ}\text{C}$) and methylmercury contamination on A) Fulton condition index and B) the encephalization coefficient of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Different letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition.

4.3. Macro areas coefficient

Ocean warming, acidification and contamination did not cause any significant changes in the macro areas coefficient ratio between telencephalon, optic tectum, diencephalon, cerebellum (Fig. 9A, B, C and D; $P > 0.05$). Brain stem was significantly bigger than what was predicted under warming and control pH (Fig. 9E; $P < 0.05$). Between non-contaminated and contaminated warming and control pH conditions there was a significant difference in the brain stem coefficient ($P < 0.05$).






Figure 9. Impact of ocean acidification ($\Delta\text{pH } 0.5$), warming ($+4\text{ }^{\circ}\text{C}$) and methylmercury contamination on macroareas coefficient between the five areas of the brain A) telencephalon, B) optic tectum, C) diencephalon, D) cerebellum and E) brain stem of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition.

4.4. Brain macro areas acetylcholinesterase activity

Ocean warming and acidification did not affect significantly AChE activity in all five meagre brain macro areas under non-contaminated conditions (Fig. 10; $P > 0.05$). However, contamination induced a significant increase in AChE activity in all areas (Fig. 10; $P < 0.05$). In the telencephalon, warming and acidification revealed an increase in AChE (1575.01 nmol min⁻¹ mg⁻¹ protein) compared to the activity under the same temperature condition and control pH (830.96 nmol min⁻¹ mg⁻¹ protein) (Fig. 10A; $P < 0.05$). Moreover, AChE in the optic tectum increased significantly under the most drastic scenario (23 °C and pH 7.5) (1434.15 nmol min⁻¹ mg⁻¹ protein) (Fig. 10B; $P < 0.05$). In diencephalon, AChE decreased under the warming and control pH contaminated treatment (962.35 nmol min⁻¹ mg⁻¹ protein) in comparison with the same temperature but hypercapnia treatment (1531.92 nmol min⁻¹ mg⁻¹ protein) (Fig. 10C; $P < 0.05$). In contrast, in the cerebellum, neither temperature nor pH affected the enzyme response to contamination (Fig. 10D, $P > 0.05$). Under the contaminated scenario, warming and hypercapnia increased AChE activity in brain stem (1837.70 nmol min⁻¹ mg⁻¹ protein) in comparison with control temperature and pH, and control temperature and hypercapnia treatments (1376.38 nmol min⁻¹ mg⁻¹ protein and 1375.70 nmol min⁻¹ mg⁻¹ protein, respectively) (Fig. 10E; $P < 0.05$).

Only the non-contaminated control temperature and pH and warming and control pH treatments showed significant differences between the five brain macro areas AChE activities under the same treatment. Under contaminated conditions, only the warming and hypercapnia condition did not reveal significant differences in the AChE levels of the different brain macro areas under the same treatment. Under the non-contaminated temperature and pH control condition, AChE was higher in the diencephalon (243.32 nmol min⁻¹ mg⁻¹ protein) than in the brain stem (141.08 nmol min⁻¹ mg⁻¹ protein) (Fig. 10; $P < 0.05$). In the non-contaminated condition, under warming and control pH, telencephalon and cerebellum had lower AChE activity (125.11 nmol min⁻¹ mg⁻¹ protein and 117.45 nmol min⁻¹ mg⁻¹ protein, respectively) than diencephalon and brain stem (206.88 nmol min⁻¹ mg⁻¹ protein and 217.61 nmol min⁻¹ mg⁻¹ protein, respectively) (Fig. 10; $P < 0.05$). Under the contaminated control temperature and pH condition, cerebellum activity was significantly lower (794.28 nmol min⁻¹ mg⁻¹ protein) than in the brain stem (1376.39 nmol min⁻¹ mg⁻¹ protein) (Fig. 10; $P < 0.05$). Under the control temperature and hypercapnia contaminated condition, diencephalon AChE activity (1475.72 nmol min⁻¹ mg⁻¹ protein) was statistically higher than in the cerebellum (931.20 nmol min⁻¹ mg⁻¹ protein) (Fig. 10; $P < 0.05$). Lastly, under the warming and control pH scenario, telencephalon, diencephalon and cerebellum had lower AChE

activity ($831.00 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$, $962.35 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$ and $801.95 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$, respectively) than brain stem ($1461.22 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$) (Fig. 10; $P < 0.05$).




Figure 10. Impact of ocean acidification (Δ pH 0.5), warming (+4 °C) and methylmercury contamination on the acetylcholinesterase activity on the five areas of the brain A) telencephalon, B) optic tectum, C) diencephalon, D) cerebellum and E) brain stem of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition. Capital letters represent significant differences between tissues in the same treatment.

4.5. Heat shock response and lipid peroxidation

Water temperature, pH and the usage of a non-contaminated/contaminated food source had distinct effects on HSP70/HSC70 expression in different tissues. Only muscle displayed significant differences (Fig. 11B; $P < 0.05$), i.e. an increase in the HSP content occurred under both warming and control pH conditions (5.55 ng mg⁻¹ protein in the non-contaminated condition and 6.06 ng mg⁻¹ protein in the contaminated condition) when compared to control conditions (19 °C pH 8.0) (3.76 ng mg⁻¹ protein in the non-contaminated condition and 3.51 ng mg⁻¹ protein in the contaminated condition). However, there was a significant drop in HSP content, under warming and hypercapnia conditions (23 °C pH 7.5) in non-contaminated tissue (3.94 ng mg⁻¹ protein) (Fig. 11B; $P < 0.05$).

When comparing different tissues in the same condition it is evident that the HSP content changes (Fig. 11). Under control non-contaminated and contaminated conditions, HSP was statistically higher in liver (5.85 ng mg⁻¹ protein in the non-contaminated condition and 6.68 ng mg⁻¹ protein in the contaminated condition) than in gills (3.47 ng mg⁻¹ protein in the non-contaminated condition and 3.89 ng mg⁻¹ protein in the contaminated condition) and muscle (3.76 ng mg⁻¹ protein in the non-contaminated condition and 3.51 ng mg⁻¹ protein in the contaminated condition) ($P < 0.05$; Fig. 11). At warming and control pH scenario, HSP's were significantly higher in the muscle (5.55 ng mg⁻¹ protein) when compared to gills and liver (3.19 ng mg⁻¹ protein and 3.60 ng mg⁻¹ protein, respectively) ($P < 0.05$; Fig. 11). Under the non-contaminated warming and hypercapnia scenario, HSP's were statistically higher in the muscle and liver (3.94 ng mg⁻¹ protein and 4.77 ng mg⁻¹ protein, respectively) ($P < 0.05$; Fig. 11).

Regarding MDA levels, no significant differences were observed between treatments ($P > 0.05$; Fig 11D).




Figure 11. Impact of ocean acidification (Δ pH 0.5), warming (+4 °C) and methylmercury contamination on HSP in A) gills, B) muscle and C) liver and on D) muscle MDA concentration in juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Small letters represent statistical differences among treatments in the same contamination condition. Capital letters represent significant differences in HSP between tissues in the same treatment.

4.6. Antioxidant enzymes activities

The impact of ocean warming, acidification and contamination on antioxidant enzymes (GST, CAT and SOD) in the muscle of juvenile *Argyrosomus regius* is shown in Figure 12.

GST activity did not change with contamination (Fig. 12A; $P > 0.05$). Warming caused an increase in GST activity ($P < 0.05$). Furthermore, under contamination, GST was statistically the highest under warming and control pH scenario when compared with warming and acidification, with values ranging from $9.47 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$ to $6.05 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$, respectively, ($P < 0.05$).

Catalase activity did not display significant differences in non-contaminated treatments (Fig.12B; $P > 0.05$), however when contaminated with MeHg, meagre CAT activity statistically increased from 4.22 to $6.64 \text{ nmol min}^{-1} \text{ mg}^{-1} \text{ protein}$ under the control temperature and hypercapnia scenario ($P < 0.05$).

SOD activity did not show any significant differences under non-contaminated treatments (Fig.12C; $P > 0.05$). In contrast, in contaminated conditions, SOD was significantly higher under the control temperature and acidification treatment ($P < 0.05$).




Figure 12. Impact of ocean acidification (Δ pH 0.5), warming (+4 °C) and methylmercury contamination in the A) GST activity, B) Catalase activity and C) SOD activity in the muscle tissue of juvenile meagre (*Argyrosomus regius*). Values represent mean+s.d. Different small letters represent statistical differences among treatments in the same contamination condition. Asterisk represents significant differences between contaminated and non-contaminated treatments in each temperature and pH condition.

5. Discussion

5.1. Total mercury accumulation

Aquatic organisms can accumulate mercury from water, through gills, or from feed ingestion (Jernelöv and Lann, 1971). Here, the contamination occurred through feed intake. Furthermore, mercury accumulation in fish depends on the water quality of their habitat (temperature, pH, alkalinity, etc...) (Harris and Bodaly, 1998; Ponce and Bloom, 1991; Wren et al., 1991). Temperature can influence mercury bioaccumulation in fish (Harris and Bodaly, 1998) with evidence from killifish to rainbow trout (Dijkstra et al., 2013; MacLeod and Pessah, 1973). Moreover, several studies have demonstrated that shifts in pH can also influence mercury accumulation in fish. For example, Scheuhammer and Graham, (1999) revealed that fish from an acidic lake accumulated more mercury compared to fish from a neutral lake, thus concluding that pH is negatively correlated with mercury concentrations in freshwater fish (Ponce and Bloom, 1991; Wren et al., 1991) and also invertebrates (Wren and Stephenson, 1991). In the present study, warming increased mercury accumulation in fish compared with the control condition, though only significantly in gills and muscle. Despite the evidence that low pH might increase mercury accumulation in freshwater fish (Haines et al., 1992; Ponce and Bloom, 1991), the current findings did not reflect this pattern. It is worth noting that possible evolutionary adaptations might influence the different results observed compared to the previous studies with freshwater.

Affinity to metal accumulation can differ between fish organs. Muscle is usually the organ with less affinity to metals (Jeziarska and Witeska, 2006) compared to liver where metals accumulate at higher levels (Jeziarska and Witeska, 2006), as the latest tissue plays a major role in accumulation and detoxification of metals (Gbem et al., 2001; Wagner and Boman, 2003). Mercury levels are usually higher in the liver also because this contaminant is absorbed at the intestine from the feed, being an important route for mercury uptake, and then transported to the liver through the hepatic portal system (Foster et al., 2000). Indeed, in this study muscle accumulated less mercury in most treatments, whereas liver had the highest mercury accumulation in most treatments. Higher concentrations of Hg have been found in liver in striped bass, catfish, and tilapia in comparison with Hg concentrations in skeletal muscle, liver, blood, gonad, brain, gill, and heart of these species (Cizdziel et al., 2003). Brain is an organ that usually also shows high metal accumulation (Jeziarska and Witeska, 2006). Gills displayed lower mercury accumulation. This tissue is very sensitive to metal accumulation when waterborne metal exposure occurs (De Domenico et al., 2011). However, in this study

contamination was made through feed intake, which may have caused gills to accumulate less mercury under the warming and acidification condition. It is known that the interaction between warming and acidification can have additive or antagonistic effects on gills mercury uptake (Byrne and Przeslawski, 2013), likely justifying the results obtained in this study.

5.2. Fulton condition and survival

In the present study, meagre did not demonstrated any changes in survival and Fulton condition under ocean warming, acidification and contamination. Nonetheless, MacLeod and Pessah, (1973) revealed that mortality rate, due to mercury exposure, is linearly related to temperature in freshwater fish. In zebrafish, survival is affected by mercury too (Penglase et al., 2014). Moreover, mercury concentrations are negatively correlated with Fulton condition in wild river fish populations (Pyle et al., 2005). Such results indicate that fish species can respond differently to environmental changes and contamination levels. In fact, meagre is a very resilient species and can easily adapt to environmental alterations (Monfort, 2010), which may explain the absence of such expected effects.

5.3. Encephalization

Teleost fish brain grows continuously throughout fish lifetime (Ekström, 1994; Kotrschal et al., 1998). Hence, due to brain plasticity this organ can be affected by diverse environmental changes that occur during fish life (Kihlslinger and Nevitt, 2006). It has been reported that brain size can be influenced by habitat complexity (Pollen et al., 2007; Shumway, 2010) and type of diet (Gonzalez-Voyer et al., 2009) in African cichlid fish. Rearing conditions can also affect brain size in Atlantic salmon, as fish reared in stream conditions can develop smaller brains compared with fish in hatcheries (Näslund et al., 2012). Kotrschal et al., (2012) also noted that animals reared in hatchery developed larger brains. In contrast, Marchetti and Nevitt, (2003) observed that brains of hatchery reared rainbow trout are smaller than wild fish. Furthermore, hypoxia can also have effects in brain size. Chapman and Hulen, (2001) showed that brain size of two mormyrids species from extremely hypoxic waters are significant smaller than the brain of two open-water species from well oxygenated habitats. Furthermore, Chapman et al., (2008) demonstrated that brain size is smaller in African cichlids raised under hypoxia.

Yu et al., (2014) formulated a framework that identifies temperature as the most critical factor to brain enlargement and shows that fish in warmer habitats have larger brain than those living in colder waters. Gillooly and McCoy, (2014) also discussed that brain size increases with temperature in vertebrates, suggesting that higher temperature

can increase energy supply by speeding biochemical reaction rates and allowing fish to keep more brain tissue. In the present study, this effect did not occur under the warming condition. However, brain was larger under the warming and acidification treatment on both non-contaminated and contaminated treatments. The interaction between warming and acidification can have additive or antagonistic effects in marine organisms (Byrne and Przeslawski, 2013; Crain et al., 2008). Thus, hypercapnia may have enhanced the effect of temperature on brain and increased its size. Meagre may have invested more energy in brain development instead of relocating such energy to other functions as a response to the synergistic condition, which could be advantageous since fish benefits with larger brains as this tissue has innumerable important functions (Crispo and Chapman, 2010). Also, larger brains may promote the development of different behavioural patterns helping fish dealing more successfully with new environmental challenges (Sol, 2009). So the brain increase observed under the warming and acidification treatment may have been an attempt of meagre to adapt to the environmental conditions.

Mercury can also cause brain lesions in fish (Berntssen et al., 2003). Encephalization coefficient was significantly different between the non-contaminated scenario (with a significantly larger brain mass) and the contaminated one, suggesting that mercury significantly affected brain investment.

5.4. Brain macro areas

Environment conditions can affect brain structures, including its size and volume (Kihlslinger et al., 2006; Kotrschal et al., 2012). Kihlslinger et al., (2006) reported that olfactory bulb and telencephalon are larger in wild fish compared to hatchery-reared fish in a single generation of Chinook salmon. Different rearing conditions also cause an increase in the cerebellar area of salmon alevin (Kihlslinger and Nevitt, 2006). Furthermore, different hatchery-rearing practices influence different brain structures like the optic tectum and the telencephalon in rainbow trout (Marchetti and Nevitt, 2003). Thus, it is possible to see different responses in brain structures to environmental stressors. In the present study, brain stem was the only brain area affected by environmental conditions. This structure controls sensory systems (except smell and sight) and governs certain somatic and visceral functions. It also controls respiration and osmoregulation in bony fish (Helfman et al., 2009).

Fish can accumulate mercury in specific areas of the brain. Rouleau et al., (1999) showed that rainbow and brown trout exposed to waterborne mercury can accumulate this metal in several brain areas, for example, in cerebellum and optic tectum. This suggests that in these fish species waterborne Hg was taken by water-exposed receptor

cells of sensory nerves such as: receptors of lateral line system, cutaneous sensory cells, and receptor cells of taste buds in oral mucosa and then transferred to the different areas of the brain through axonal transport (a physiological process for the transport of dissolved neuronal constituents along nerve axons). In this study, contamination was made through feed intake, which can lead to an uptake of mercury through taste buds and subsequently transported to the brain. However, contamination did not affect brain areas in the present study, although brain have shown high levels of mercury contamination, probably due to the fact that meagre is a resilient species and easily adaptable to different environmental conditions (Monfort, 2010).

As previously said, high temperature is linked to larger brains (Gillooly and McCoy, 2014; Yu et al., 2014). In this study, warming affected brain stem size, and caused an enlargement in this structure, likely as a response to the natural increase in metabolic rates, energy supply, and biochemical reaction rates that occur at higher temperatures (Gillooly and McCoy, 2014), favouring an increased investment in brain stem.

5.5. Acetylcholinesterase (AChE) activity

Diverse AChE responses have been reported when exposed to different toxic compounds (Richetti et al., 2011; Romani et al., 2003). Although temperature can alter AChE activity (Baldwin, 1971; Hogan, 1971), future temperature conditions did not affect AChE response in meagre. It has been described that ocean acidification can alter GABA function in fish brain (Nilsson et al., 2012). GABA is an important neurotransmitter in the central nervous system (CNS) as well as acetylcholine (Ach), which is hydrolyzed by AChE. Hence, it is expected that acidification alter AChE activity in meagre. However, this was not observed. Meagre is a high resilient species, which may explain why warming and acidification under non-contaminated treatments did not affect significantly AChE activity.

Here, it is also shown that AChE activity in the five brain areas was remarkably higher in contaminated meagre. In contrast, several studies have shown a decrease in fish AChE in the presence of mercury and other metals (Gill et al., 1990; Richetti et al., 2011). AChE also seems to have different responses to other contaminants. In fact, an increase in AChE activity has been demonstrated in fish exposed to copper, zinc and ethanol (Dethloff et al., 1999; Rico et al., 2007; Romani et al., 2003).

Interestingly, in vertebrates it has already been reported that the exposure to MeHg compromise the cerebral dopaminergic and serotonergic systems according to the brain area (Castoldi et al., 2006). This suggests that the neurotransmitter Ach may also be affected by MeHg, likely explaining the variation in AChE activities among the

different brain macro areas for the same treatment, as each area can have a distinct way to respond to environmental stresses. Except for cerebellum and brain stem, under contaminated treatments, the warming and acidification condition showed higher AChE activity than the warming treatment. The synergism between the two factors probably exacerbated the effect of contamination, enhancing AChE activity in meagre brain. Moreover, this synergism has also been demonstrated to increase AChE levels in sharks (Rosa et al., 2015 under submission). In optic tectum and brain stem a difference between the effect of acidification and the synergistic condition (high temperature and low pH) was observed. As expected, under this condition, AChE levels were higher, showing once again that synergism aggravated the effect of contamination.

It is worth noting that AChE can be involved in the regulation of apoptosis (Jin et al., 2004; Zhang et al., 2002). Since mercury can induce apoptosis in fish (Drevnick et al., 2006; Sarmiento et al., 2004), the increase in AChE activity in contaminated treatments likely indicate the initiation of this mechanism (De Domenico et al., 2013; Jin et al., 2004; Zhang et al., 2002). Furthermore, oxidative stress increases AChE activity (Melo et al., 2003) and Hg is responsible for oxidative stress in fish brain (Berntssen et al., 2003). Therefore, Hg may have triggered oxidative stress in meagre brain, leading to an increase in AChE activity.

5.6. Oxidative stress

High temperatures might lead to the formation of reactive oxygen species (ROS) (Abele et al., 2002; Lesser, 2006). HSP are one of the molecules that protect and control ROS formation (Jacquier-Sarlin et al., 1994). These proteins help to repair, refold and eliminate damaged or denatured proteins (Sokolova et al., 2011). In non-contaminated and contaminated warming conditions, HSP in muscle showed an increase, demonstrating a stress response in fish. Acidification did not change HSP response indicating that the effect of temperature on HSP is stronger than acidification. Under the non-contaminated condition, the synergism between high temperature and low pH diminished the heat effect on HSP, demonstrating that acidification exhibit an antagonistic effect.

In muscle, contamination did not affect HSP response. Gills are usually the most affected tissues when fish is exposed to contamination, particularly when contamination occurs through water (Farag et al., 1994). In the present study, the contamination occurred via diet, and therefore gills were less affected (revealing the smallest HSP response) when compared to liver and muscle. Liver did not exhibit any differences in HSP response among the treatments, despite displaying the highest levels of HSP in

comparison with other tissues, likely since liver plays a major role in the detoxification of xenobiotics (Williams et al., 1996).

Lipid peroxidation is one of the most common cellular injury mechanisms (Lesser, 2006). MDA concentration has not changed in any treatment, indicating that ROS did not have an effect on meagre lipids (peroxidation). Thus, it is possible to conclude that HSP response and antioxidant enzymes activities seemed to work efficiently against cellular injuries.

Antioxidant enzymes are another defense mechanism against oxidative stress. Among them, GST is involved in the protection against xenobiotics (Wang et al., 2000), SOD converts superoxide (O_2^-) into hydrogen peroxide (H_2O_2) and CAT converts H_2O_2 into water (H_2O) and oxygen (O_2) (Lesser, 2006). Antioxidant enzymes generally increase in response to ROS production, which in turn is increased with warming (Abele et al., 2002; Lesser, 2006). It has been reported that GST increases in response to warming in marine animals, such as sole post-metamorphic larvae (Pimentel et al., 2015) and tropical and temperate shrimp (Rosa et al., 2014a). Hence, as expected, meagre GST increased at 23 °C, in order to regulate ROS production. Similarly to HSP, significant differences were only observed in the warming treatment, showing once more that warming has a more drastic effect on meagre than acidification. The synergism between warming and acidification decrease GST activity in the contaminated treatment, indicating that once again acidification has an antagonistic effect in temperature, decreasing its negative effect in GST activity.

Catalase generally increased in response to contamination. However this effect was only significant under the acidification and control temperature treatment. Previous studies reported contradictory responses of catalase to metal contamination. For instance, silver decrease CAT activity, whereas cadmium, zinc and chromium increased CAT activity in the liver of freshwater fish (Atli et al., 2006). In another study with the same fish model, liver CAT activity increased with cadmium and lead exposures, but was inhibited by zinc (Atli and Canli, 2007). Furthermore, Mieiro et al., (2011) observed CAT depletion in mercury contaminated meagre. In the present study, CAT activity increased in samples from contaminated treatments, likely to control ROS production. Also, the absence of significant differences between the various non-contaminated treatments in CAT activity may also suggest that meagre is a tolerant species to high temperatures and acidification.

In the present study, SOD did not exhibit significant differences between non-contaminated and contaminated treatments. This enzyme activity only revealed differences between contaminated treatments, e.g. acidification increased SOD activity. It is not well studied if hypercapnia causes ROS increase in fish. However, Tomanek et

al., (2011) reported that hypercapnia may enhance oxidative stress in oysters. Although acidification did not have an effect on GST, SOD activity increased under the low pH contaminated treatment possibly to protect fish from ROS. This enzyme can also increase in response to MeHg, suggesting an adaptive response of the redox defense system (Berntssen et al., 2003). So, contamination and acidification may have triggered a defense response in meagre, causing the increase in SOD activity.

6. Final remarks

The present study shows that meagre will not suffer major consequences under the expected future ocean contaminated conditions. Nonetheless, here it is shown that warming will be the most relevant factor that will increase mercury bioaccumulation in meagre. Some variables were only affected in contaminated treatments, especially AChE. Although meagre suffered some physiological alterations under these conditions, meagre responses were not apparently harmful for fish. These results suggest that meagre will be able to adapt to the predicted future ocean conditions even under mercury contamination. However, future studies will be necessary to comprehensively understand how marine biota will respond to the combined effects of future warming, acidification and pollution.

References

- Abele D, Heise K, Pörtner H-O, Puntarulo S. Temperature-dependence of mitochondrial function and production of reactive oxygen species in the intertidal mud clam *Mya arenaria*. *Journal of Experimental Biology*. 2002; 205:1831-41.
- Atli G, Alptekin Ö, Tükel S, Canli M. Response of catalase activity to Ag⁺, Cd²⁺, Cr⁶⁺, Cu²⁺ and Zn²⁺ in five tissues of freshwater fish *Oreochromis niloticus*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2006; 143:218-24.
- Atli G, Canli M. Enzymatic responses to metal exposures in a freshwater fish *Oreochromis niloticus*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2007; 145:282-7.
- Azevedo LB, De Schryver AM, Hendriks AJ, Huijbregts MA. Calcifying species sensitivity distributions for ocean acidification. *Environmental Science & Technology*. 2015; 49:1495-500.
- Baatrup E. Structural and functional effects of heavy metals on the nervous system, including sense organs, of fish. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology*. 1991; 100:253-7.
- Baldwin J. Adaptation of enzymes to temperature: acetylcholinesterases in the central nervous system of fishes. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*. 1971; 40:181-7.
- Berntssen MH, Aatland A, Handy RD. Chronic dietary mercury exposure causes oxidative stress, brain lesions, and altered behaviour in Atlantic salmon (*Salmo salar*) parr. *Aquatic Toxicology*. 2003; 65:55-72.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*. 1976; 72:248-54.
- Burger J, Dixon C, Boring S, Gochfeld M. Effect of deep-frying fish on risk from mercury. *Journal of Toxicology and Environmental Health Part A*. 2003; 66:817-28.
- Byrne M, Przeslawski R. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology*. 2013; 53:582-96.
- Caldeira K, Wickett ME. Oceanography: anthropogenic carbon and ocean pH. *Nature*. 2003; 425:365.
- Campbell AL, Mangan S, Ellis RP, Lewis C. Ocean Acidification Increases Copper Toxicity to the Early Life History Stages of the Polychaete *Arenicola marina* in Artificial Seawater. *Environmental Science & Technology*. 2014; 48:9745-53.

- Campbell LM, Norstrom RJ, Hobson KA, Muir DC, Backus S, Fisk AT. Mercury and other trace elements in a pelagic Arctic marine food web (Northwater Polynya, Baffin Bay). *Science of the Total Environment*. 2005; 351:247-63.
- Castoldi AF, Blandini F, Randine G, Samuele A, Manzo L, Coccini T. Brain monoaminergic neurotransmission parameters in weanling rats after perinatal exposure to methylmercury and 2, 2', 4, 4', 5, 5'-hexachlorobiphenyl (PCB153). *Brain Research*. 2006; 1112:91-8.
- Chapman L, Albert J, Galis F. Developmental plasticity, genetic differentiation, and hypoxia-induced trade-offs in an African cichlid fish. *The Open Evolution Journal*. 2008; 2:75-88.
- Chapman LJ, Hulén KG. Implications of hypoxia for the brain size and gill morphometry of mormyrid fishes. *Journal of Zoology*. 2001; 254:461-72.
- Checkley DM, Dickson AG, Takahashi M, Radich JA, Eisenkolb N, Asch R. Elevated CO₂ enhances otolith growth in young fish. *Science*. 2009; 324:1683-.
- Cizdziel J, Hinnert T, Cross C, Pollard J. Distribution of mercury in the tissues of five species of freshwater fish from Lake Mead, USA. *Journal of Environmental Monitoring*. 2003; 5:802-7.
- Crain CM, Kroeker K, Halpern BS. Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology letters*. 2008; 11:1304-15.
- Cripps IL, Munday PL, McCormick MI. Ocean acidification affects prey detection by a predatory reef fish. *PLOS ONE*. 2011; 6:e22736.
- Crispo E, Chapman L. Geographic variation in phenotypic plasticity in response to dissolved oxygen in an African cichlid fish. *Journal of Evolutionary Biology*. 2010; 23:2091-103.
- De Domenico E, Mauceri A, Giordano D, Maisano M, Giannetto A, Parrino V, et al. Biological responses of juvenile European sea bass (*Dicentrarchus labrax*) exposed to contaminated sediments. *Ecotoxicology and Environmental Safety*. 2013; 97:114-23.
- De Domenico E, Mauceri A, Giordano D, Maisano M, Giofrè G, Natalotto A, et al. Effects of "in vivo" exposure to toxic sediments on juveniles of sea bass (*Dicentrarchus labrax*). *Aquatic Toxicology*. 2011; 105:688-97.
- Dethloff GM, Schlenk D, Hamm JT, Bailey HC. Alterations in physiological parameters of rainbow trout (*Oncorhynchus mykiss*) with exposure to copper and copper/zinc mixtures. *Ecotoxicology and Environmental Safety*. 1999; 42:253-64.
- Dickson A, Millero F. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research Part A Oceanographic Research Papers*. 1987; 34:1733-43.

- Dijkstra JA, Buckman KL, Ward D, Evans DW, Dionne M, Chen CY. Experimental and Natural Warming Elevates Mercury Concentrations in Estuarine Fish. PLOS ONE. 2013; 8
- Dorts J, Kestemont P, Thézenas M-L, Raes M, Silvestre F. Effects of cadmium exposure on the gill proteome of *Cottus gobio*: Modulatory effects of prior thermal acclimation. Aquatic Toxicology. 2014; 154:87-96.
- Drevnick PE, Sandheinrich MB, Oris JT. Increased ovarian follicular apoptosis in fathead minnows (*Pimephales promelas*) exposed to dietary methylmercury. Aquatic Toxicology. 2006; 79:49-54.
- Ekström P. Developmental changes in the brain-stem serotonergic nuclei of teleost fish and neural plasticity. Cellular and Molecular Neurobiology. 1994; 14:381-93.
- El-Shebly AA, El-Kady MA, Hussin A, Hossain MY. Preliminary observations on the pond culture of meagre, *Argyrosomus regius* (Asso, 1801)(Sciaenidae) in Egypt. Journal of Fisheries and Aquatic Science. 2007; 2:345-52.
- Ellman GL, Courtney KD, Andres V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochemical Pharmacology. 1961; 7:88-95.
- EPA. Method 7473 Mercury in solids and solutions by thermal decomposition, amalgamation, and atomic absorption spectrophotometry: U.S. Environmental Protection Agency; 2007.
- Evers DC, Wiener JG, Basu N, Bodaly R, Morrison HA, Williams KA. Mercury in the Great Lakes region: bioaccumulation, spatiotemporal patterns, ecological risks, and policy. Ecotoxicology. 2011; 20:1487-99.
- Farag AM, Boese CJ, Bergman HL, Woodward D. Physiological changes and tissue metal accumulation in rainbow trout exposed to foodborne and waterborne metals. Environmental Toxicology and Chemistry. 1994; 13:2021-9.
- Foster E, Drake D, DiDomenico G. Seasonal changes and tissue distribution of mercury in largemouth bass (*Micropterus salmoides*) from Dorena Reservoir, Oregon. Archives of Environmental Contamination and Toxicology. 2000; 38:78-82.
- Gbem T, Balogun J, Lawal F, Annune P. Trace metal accumulation in *Clarias gariepinus* (Teugels) exposed to sublethal levels of tannery effluent. Science of the Total Environment. 2001; 271:1-9.
- Gill TS, Tewari H, Pande J. Use of the fish enzyme system in monitoring water quality: effects of mercury on tissue enzymes. Comparative Biochemistry and Physiology Part C: Comparative Pharmacology. 1990; 97:287-92.
- Gillooly JF, McCoy MW. Brain size varies with temperature in vertebrates. PeerJ. 2014; 2:e301.

- Gonçalves D, Teles M, Alpedrinha J, Oliveira RF. Brain and gonadal aromatase activity and steroid hormone levels in female and polymorphic males of the peacock blenny *Salarias pavo*. *Hormones and Behavior*. 2008; 54:717-25.
- Gonzalez-Voyer A, Winberg S, Kolm N. Social fishes and single mothers: brain evolution in African cichlids. *Proceedings of the Royal Society of London B: Biological Sciences*. 2009; 276:161-7.
- Gonzalez P, Dominique Y, Massabuau J, Boudou A, Bourdineaud J. Comparative effects of dietary methylmercury on gene expression in liver, skeletal muscle, and brain of the zebrafish (*Danio rerio*). *Environmental Science & Technology*. 2005; 39:3972-80.
- Gray JE, Theodorakos PM, Bailey EA, Turner RR. Distribution, speciation, and transport of mercury in stream-sediment, stream-water, and fish collected near abandoned mercury mines in southwestern Alaska, USA. *Science of the Total Environment*. 2000; 260:21-33.
- Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *Journal of Biological Chemistry*. 1974; 249:7130-9.
- Haines TA, Komov V, Jagoe CH. Lake acidity and mercury content of fish in Darwin National Reserve, Russia. *Environmental Pollution*. 1992; 78:107-12.
- Hammerschmidt CR, Sandheinrich MB, Wiener JG, Rada RG. Effects of dietary methylmercury on reproduction of fathead minnows. *Environmental Science & Technology*. 2002; 36:877-83.
- Han Z-X, Wu D-D, Wu J, Lv C-X, Liu Y-R. Effects of Ocean Acidification on Toxicity of Heavy Metals in the Bivalve *Mytilus edulis* L. *Synthesis and Reactivity in Inorganic Metal-Organic and Nano-Metal Chemistry*. 2014; 44:133-9.
- Harley CD, Randall Hughes A, Hultgren KM, Miner BG, Sorte CJ, Thornber CS, et al. The impacts of climate change in coastal marine systems. *Ecology Letters*. 2006; 9:228-41.
- Harris RC, Bodaly RD. Temperature, growth and dietary effects on fish mercury dynamics in two Ontario lakes. *Biogeochemistry*. 1998; 40:175-87.
- Harvey BP, Gwynn-Jones D, Moore PJ. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution*. 2013; 3:1016-30.
- Helfman G, Collette BB, Facey DE, Bowen BW. *The diversity of fishes: biology, evolution, and ecology*: John Wiley & Sons; 2009. 720 pp
- Hoegh-Guldberg O, Mumby P, Hooten A, Steneck R, Greenfield P, Gomez E, et al. Coral reefs under rapid climate change and ocean acidification. *Science*. 2007; 318:1737-42.

- Hogan JW. Water temperature as a source of variation in specific activity of brain acetylcholinesterase of bluegills. *Bulletin of Environmental Contamination and Toxicology*. 1971; 5:347-53.
- IPCC. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2013. 1535 pp
- IPCC. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2014. 1132 pp
- Ishimatsu A, Hayashi M, Kikkawa T. Fishes in high-CO₂, acidified oceans. *Marine Ecology Progress Series*. 2008; 373:295-302.
- Jacquier-Sarlin M, Fuller K, Dinh-Xuan A, Richard M-J, Polla B. Protective effects of hsp70 in inflammation. *Experientia*. 1994; 50:1031-8.
- Jernelöv A, Lann H. Mercury accumulation in food chains. *Oikos*. 1971:403-6.
- Jezierska B, Witeska M. The metal uptake and accumulation in fish living in polluted waters. *Soil and water pollution monitoring, protection and remediation*: Springer; 2006. 107-14
- Jin Q, He H, Shi Y, Lu H, Zhang X. Overexpression of acetylcholinesterase inhibited cell proliferation and promoted apoptosis in NRK cells. *Acta Pharmacologica Sinica*. 2004; 25:1013-21.
- Johansson LH, Borg LH. A spectrophotometric method for determination of catalase activity in small tissue samples. *Analytical Biochemistry*. 1988; 174:331-6.
- Keyvanshokoo S, Vaziri B, Gharaei A, Mahboudi F, Esmaili-Sari A, Shahriari-Moghadam M. Proteome modifications of juvenile beluga (*Huso huso*) brain as an effect of dietary methylmercury. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*. 2009; 4:243-8.
- Kihlslinger R, Lema SC, Nevitt G. Environmental rearing conditions produce forebrain differences in wild Chinook salmon *Oncorhynchus tshawytscha*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2006; 145:145-51.
- Kihlslinger RL, Nevitt GA. Early rearing environment impacts cerebellar growth in juvenile salmon. *Journal of Experimental Biology*. 2006; 209:504-9.
- Korbas M, MacDonald TC, Pickering IJ, George GN, Krone PH. Chemical form matters: differential accumulation of mercury following inorganic and organic mercury exposures in zebrafish larvae. *ACS Chemical Biology*. 2011; 7:411-20.

- Kotrschal A, Sundström LF, Brelin D, Devlin R, Kolm N. Inside the heads of David and Goliath: environmental effects on brain morphology among wild and growth-enhanced coho salmon *Oncorhynchus kisutch*. *Journal of Fish Biology*. 2012; 81:987-1002.
- Kotrschal K, Van Staaden M, Huber R. Fish brains: Evolution and environmental relationships. *Reviews in Fish Biology and Fisheries*. 1998; 8:373-408.
- Lacoue-Labarthe T, Martin S, Oberhänsli F, Teyssié J-L, Jeffree R, Gattuso J-P, et al. Temperature and $p\text{CO}_2$ effect on the bioaccumulation of radionuclides and trace elements in the eggs of the common cuttlefish, *Sepia officinalis*. *Journal of Experimental Marine Biology and Ecology*. 2012; 413:45-9.
- Lacoue-Labarthe T, Martin S, Oberhänsli F, Teyssié J-L, Markich S, Jeffree R, et al. Effects of increased $p\text{CO}_2$ and temperature on trace element (Ag, Cd and Zn) bioaccumulation in the eggs of the common cuttlefish, *Sepia officinalis*. *Biogeosciences*. 2009; 6:2561-73.
- Lacoue-Labarthe T, Reveillac E, Oberhänsli F, Teyssié J-L, Jeffree R, Gattuso J. Effects of ocean acidification on trace element accumulation in the early-life stages of squid *Loligo vulgaris*. *Aquatic Toxicology*. 2011; 105:166-76.
- Lapointe D, Pierron F, Couture P. Individual and combined effects of heat stress and aqueous or dietary copper exposure in fathead minnows (*Pimephales promelas*). *Aquatic Toxicology*. 2011; 104:80-5.
- Larose C, Canuel R, Lucotte M, Di Giulio RT. Toxicological effects of methylmercury on walleye (*Sander vitreus*) and perch (*Perca flavescens*) from lakes of the boreal forest. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2008; 147:139-49.
- Lesser MP. Oxidative stress in marine environments: biochemistry and physiological ecology. *Annual Review of Physiology*. 2006; 68:253-78.
- Lewis E, Wallace D, Allison LJ. Program developed for CO_2 system calculations: Carbon Dioxide Information Analysis Center, managed by Lockheed Martin Energy Research Corporation for the US Department of Energy Tennessee; 1998.
- MacLeod J, Pessah E. Temperature effects on mercury accumulation, toxicity, and metabolic rate in rainbow trout (*Salmo gairdneri*). *Journal of the Fisheries Board of Canada*. 1973; 30:485-92.
- Magnotti RA, Eberly J, Quarm D, McConnell R. Measurement of acetylcholinesterase in erythrocytes in the field. *Clinical Chemistry*. 1987; 33:1731-5.
- Marchetti MP, Nevitt GA. Effects of hatchery rearing on brain structures of rainbow trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes*. 2003; 66:9-14.

- Matta MB, Linse J, Cairncross C, Francendese L, Kocan RM. Reproductive and transgenerational effects of methylmercury or Aroclor 1268 on *Fundulus heteroclitus*. *Environmental Toxicology and Chemistry*. 2001; 20:327-35.
- Mehrbach C, Culbertson C, Hawley J, Pytkowicz R. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*. 1973; 18:897-907.
- Melo JB, Agostinho P, Oliveira CR. Involvement of oxidative stress in the enhancement of acetylcholinesterase activity induced by amyloid beta-peptide. *Neuroscience Research*. 2003; 45:117-27.
- Mieiro C, Ahmad I, Pereira M, Duarte A, Pacheco M. Antioxidant system breakdown in brain of feral golden grey mullet (*Liza aurata*) as an effect of mercury exposure. *Ecotoxicology*. 2010; 19:1034-45.
- Mieiro C, Pereira M, Duarte A, Pacheco M. Brain as a critical target of mercury in environmentally exposed fish (*Dicentrarchus labrax*)—bioaccumulation and oxidative stress profiles. *Aquatic Toxicology*. 2011; 103:233-40.
- Monfort MC. Present market situation and prospects of meagre (*Argyrosomus regius*), as an emerging species in Mediterranean aquaculture: FAO; 2010.
- Munday PL, Crawley NE, Nilsson GE. Interacting effects of elevated temperature and ocean acidification on the aerobic performance of coral reef fishes. *Marine Ecology Progress Series*. 2009; 388:235-42.
- Munday PL, Dixon DL, McCormick MI, Meekan M, Ferrari MC, Chivers DP. Replenishment of fish populations is threatened by ocean acidification. *Proceedings of the National Academy of Sciences*. 2010; 107:12930-4.
- Näslund J, Aarestrup K, Thomassen ST, Johnsson JI, Post J. Early enrichment effects on brain development in hatchery-reared Atlantic salmon (*Salmo salar*): no evidence for a critical period. *Canadian Journal of Fisheries and Aquatic Sciences*. 2012; 69:1481-90.
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Sørensen C, Watson S-A, et al. Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nature Climate Change*. 2012; 2:201-4.
- Njemini R, Demanet C, Mets T. Comparison of two ELISAs for the determination of Hsp70 in serum. *Journal of Immunological Methods*. 2005; 306:176-82.
- Noyes PD, McElwee MK, Miller HD, Clark BW, Van Tiem LA, Walcott KC, et al. The toxicology of climate change: Environmental contaminants in a warming world. *Environment International*. 2009; 35:971-86.
- Nriagu JO, Pacyna JM. Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature*. 1988; 333:134-9.

- Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, Feely RA, et al. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*. 2005; 437:681-6.
- Penglas S, Hamre K, Ellingsen S. Selenium and mercury have a synergistic negative effect on fish reproduction. *Aquatic Toxicology*. 2014; 149:16-24.
- Pimentel M, Pegado M, Repolho T, Rosa R. Impact of ocean acidification in the metabolism and swimming behavior of the dolphinfish (*Coryphaena hippurus*) early larvae. *Marine Biology*. 2014a; 161:725-9.
- Pimentel MS, Faleiro F, Diniz M, Machado J, Pousão-Ferreira P, Peck MA, et al. Oxidative stress and digestive enzymatic activity of flatfish larvae in a changing ocean. *PLOS ONE*. 2015; 10:e0134082.
- Pimentel MS, Faleiro F, Dionísio G, Repolho T, Pousão-Ferreira P, Machado J, et al. Defective skeletogenesis and oversized otoliths in fish early stages in a changing ocean. *The Journal of Experimental Biology*. 2014b; 217:2062-70.
- Pollen AA, Dobberfuhl AP, Scace J, Igulu MM, Renn SC, Shumway CA, et al. Environmental complexity and social organization sculpt the brain in Lake Tanganyikan cichlid fish. *Brain, Behavior and Evolution*. 2007; 70:21-39.
- Ponce RA, Bloom N. Effect of pH on the bioaccumulation of low level, dissolved methylmercury by rainbow trout (*Oncorhynchus mykiss*). *Water Air & Soil Pollution*. 1991; 56:631-40.
- Pörtner H-O, Farrell AP. Physiology and climate change. *Science*. 2008; 322:690-2.
- Pyle GG, Rajotte JW, Couture P. Effects of industrial metals on wild fish populations along a metal contamination gradient. *Ecotoxicology and Environmental Safety*. 2005; 61:287-312.
- Rajeshkumar S, Munuswamy N. Impact of metals on histopathology and expression of HSP 70 in different tissues of Milk fish (*Chanos chanos*) of Kaattuppalli Island, South East Coast, India. *Chemosphere*. 2011; 83:415-21.
- Raven J, Caldeira K, Elderfield H, Hoegh-Guldberg O, Liss P, Riebesell U, et al. Ocean acidification due to increasing atmospheric carbon dioxide: The Royal Society; 2005.
- Repolho T, Baptista M, Pimentel MS, Dionísio G, Trübenbach K, Lopes VM, et al. Developmental and physiological challenges of octopus (*Octopus vulgaris*) early life stages under ocean warming. *Journal of Comparative Physiology B*. 2014; 184:55-64.
- Richetti SK, Rosemberg DB, Ventura-Lima J, Monserrat JM, Bogo MR, Bonan CD. Acetylcholinesterase activity and antioxidant capacity of zebrafish brain is altered by heavy metal exposure. *Neurotoxicology*. 2011; 32:116-22.

- Richter CA, Martyniuk CJ, Annis ML, Brumbaugh WG, Chasar LC, Denslow ND, et al. Methylmercury-induced changes in gene transcription associated with neuroendocrine disruption in largemouth bass (*Micropterus salmoides*). *General and Comparative Endocrinology*. 2014; 203:215-24.
- Rico EP, Rosemberg DB, Dias RD, Bogo MR, Bonan CD. Ethanol alters acetylcholinesterase activity and gene expression in zebrafish brain. *Toxicology Letters*. 2007; 174:25-30.
- Romani R, Antognelli C, Baldracchini F, De Santis A, Isani G, Giovannini E, et al. Increased acetylcholinesterase activities in specimens of *Sparus auratus* exposed to sublethal copper concentrations. *Chemico-Biological Interactions*. 2003; 145:321-9.
- Rosa R, Lopes AR, Pimentel M, Faleiro F, Baptista M, Trübenbach K, et al. Ocean cleaning stations under a changing climate: biological responses of tropical and temperate fish-cleaner shrimp to global warming. *Global Change Biology*. 2014a; 20:3068-79.
- Rosa R, Pimentel MS, Boavida-Portugal J, Teixeira T, Trübenbach K, Diniz M. Ocean warming enhances malformations, premature hatching, metabolic suppression and oxidative stress in the early life stages of a keystone squid. *PLOS ONE*. 2012; 7:e38282-e.
- Rosa R, Trübenbach K, Pimentel MS, Boavida-Portugal J, Faleiro F, Baptista M, et al. Differential impacts of ocean acidification and warming on winter and summer progeny of a coastal squid (*Loligo vulgaris*). *The Journal of Experimental Biology*. 2014b; 217:518-25.
- Rosa R, Trübenbach K, Repolho T, Pimentel M, Faleiro F, Boavida-Portugal J, et al. Lower hypoxia thresholds of cuttlefish early life stages living in a warm acidified ocean. *Proceedings of the Royal Society B: Biological Sciences*. 2013; 280:20131695.
- Rouleau C, Borg-Neczak K, Gottofrey J, Tjälve H. Accumulation of waterborne mercury (II) in specific areas of fish brain. *Environmental Science & Technology*. 1999; 33:3384-9.
- Rummer JL, Stecyk JA, Couturier CS, Watson S-A, Nilsson GE, Munday PL. Elevated CO₂ enhances aerobic scope of a coral reef fish. *Conservation Physiology*. 2013; 1:cot023.
- Sappal R, MacDonald N, Fast M, Stevens D, Kibenge F, Siah A, et al. Interactions of copper and thermal stress on mitochondrial bioenergetics in rainbow trout, *Oncorhynchus mykiss*. *Aquatic Toxicology*. 2014; 157:10-20.

- Sarazin G, Michard G, Prevot F. A rapid and accurate spectroscopic method for alkalinity measurements in sea water samples. *Water Research*. 1999; 33:290-4.
- Sarmiento A, Guilhermino L, Afonso A. Mercury chloride effects on the function and cellular integrity of sea bass (*Dicentrarchus labrax*) head kidney macrophages. *Fish & Shellfish Immunology*. 2004; 17:489-98.
- Scerbo R, Barghigiani C. Organic mercury determination in fish samples using an automatic mercury analyser. *Environmental Technology*. 1998; 19:339-42.
- Scheuhammer A, Graham J. The bioaccumulation of mercury in aquatic organisms from two similar lakes with differing pH. *Ecotoxicology*. 1999; 8:49-56.
- Selin NE, Jacob DJ, Yantosca RM, Strode S, Jaegle L, Sunderland EM. Global 3-D land-ocean-atmosphere model for mercury: Present-day versus preindustrial cycles and anthropogenic enrichment factors for deposition. *Global Biogeochemical Cycles*. 2008; 22
- Shumway CA. The evolution of complex brains and behaviors in African cichlid fishes. *Current Zoology*. 2010; 56:144-56.
- Sokolova IM, Lannig G. Interactive effects of metal pollution and temperature on metabolism in aquatic ectotherms: implications of global climate change. *Climate research*. 2008; 37:181.
- Sokolova IM, Sukhotin AA, Lannig G. Stress effects on metabolism and energy budgets in mollusks. *Oxidative Stress in Aquatic Ecosystems*. 2011:261-80.
- Sol D. Revisiting the cognitive buffer hypothesis for the evolution of large brains. *Biology Letters*. 2009; 5:130-3.
- Streets DG, Zhang Q, Wu Y. Projections of global mercury emissions in 2050. *Environmental Science & Technology*. 2009; 43:2983-8.
- Sun Y, Oberley LW, Li Y. A simple method for clinical assay of superoxide dismutase. *Clinical Chemistry*. 1988; 34:497-500.
- Sunderland EM, Mason RP. Human impacts on open ocean mercury concentrations. *Global Biogeochemical Cycles*. 2007; 21
- Tomanek L, Zuzow MJ, Ivanina AV, Beniash E, Sokolova IM. Proteomic response to elevated PCO₂ level in eastern oysters, *Crassostrea virginica*: evidence for oxidative stress. *The Journal of Experimental Biology*. 2011; 214:1836-44.
- Uchiyama M, Mihara M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Analytical Biochemistry*. 1978; 86:271-8.
- Wagner A, Boman J. Biomonitoring of trace elements in muscle and liver tissue of freshwater fish. *Spectrochimica Acta Part B: Atomic Spectroscopy*. 2003; 58:2215-26.

- Wang L, Groves MJ, Hepburn MD, Bowen DT. Glutathione S-transferase enzyme expression in hematopoietic cell lines implies a differential protective role for T1 and A1 isoenzymes in erythroid and for M1 in lymphoid lineages. *Haematologica*. 2000; 85:573-9.
- Weis JS, Weis P. Swimming performance and predator avoidance by mummichog (*Fundulus heteroclitus*) larvae after embryonic or larval exposure to methylmercury. *Canadian Journal of Fisheries and Aquatic Sciences*. 1995; 52:2168-73.
- Williams JH, Petersen NS, Young PA, Stansbury MA, Farag AM, Bergman HL. Accumulation of hsp70 in juvenile and adult rainbow trout gill exposed to metal-contaminated water and/or diet. *Environmental Toxicology and Chemistry*. 1996; 15:1324-8.
- Wren C, Scheider W, Wales D, Muncaster B, Gray I. Relation between mercury concentrations in walleye (*Stizostedion vitreum vitreum*) and northern pike (*Esox lucius*) in Ontario lakes and influence of environmental factors. *Canadian Journal of Fisheries and Aquatic Sciences*. 1991; 48:132-9.
- Wren C, Stephenson G. The effect of acidification on the accumulation and toxicity of metals to freshwater invertebrates. *Environmental Pollution*. 1991; 71:205-41.
- Yu Y, Karbowski J, Sachdev RN, Feng J. Effect of temperature and glia in brain size enlargement and origin of allometric body-brain size scaling in vertebrates. *BMC Evolutionary Biology*. 2014; 14:178.
- Zhang X, Yang L, Zhao Q, Caen J, He H, Jin Q, et al. Induction of acetylcholinesterase expression during apoptosis in various cell types. *Cell Death & Differentiation*. 2002; 9:790-800.
- Zhou T, Weis J. Swimming behavior and predator avoidance in three populations of *Fundulus heteroclitus* larvae after embryonic and/or larval exposure to methylmercury. *Aquatic Toxicology*. 1998; 43:131-48.

Supplemental material

Table S1. Seawater carbonate chemistry data for the different climate change and contamination scenarios.

Non-contaminated												
	19 °C						23 °C					
	8.0			7.5			8.0			7.5		
<i>measured</i>												
Temperature (°C)	19.4	±	0.4	19.0	±	0.4	23.5	±	0.1	23.3	±	0.1
Salinity	35.0	±	0.6	35.0	±	0.3	35.0	±	0.5	35.0	±	0.4
pH _T	7.99	±	0.02	7.51	±	0.19	7.97	±	0.04	7.49	±	0.11
A _T (μmol kg ⁻¹ SW)	2002.10	±	94.81	2176.81	±	71.07	1971.85	±	66.04	2093.13	±	53.90
<i>calculated</i>												
pCO ₂ (ppm)	408.46	±	20.10	1512.40	±	50.02	417.2	±	14.56	1527.82	±	39.93
C _T (μmol kg ⁻¹ SW)	1872.45	±	170.67	2139.46	±	70.79	1829.16	±	175.05	2098.78	±	119.37
Ω Ar	2.27	±	0.21	0.88	±	0.03	2.52	±	0.24	1.00	±	0.06
Contaminated												
	19 °C						23 °C					
	8.0			7.5			8.0			7.5		
<i>measured</i>												
Temperature (°C)	19.1	±	0.4	19.0	±	0.4	23.0	±	0.2	23.3	±	0.2
Salinity	34.8	±	0.4	35.0	±	0.3	35.0	±	0.5	35.1	±	0.5
pH _T	7.97	±	0.06	7.53	±	0.13	8.00	±	0.04	7.52	±	0.29
A _T (μmol kg ⁻¹ SW)	1995.02	±	83.55	2329.40	±	92.52	2046.84	±	54.98	2103.69	±	90.60
<i>calculated</i>												
pCO ₂ (ppm)	421.92	±	18.34	1532.54	±	61.65	402.67	±	11.29	1450.64	±	63.45
C _T (μmol kg ⁻¹ SW)	1850.86	±	123.61	2283.66	±	91.88	1849.83	±	98.86	2045.80	±	89.51
Ω Ar	2.15	±	0.14	0.99	±	0.04	2.65	±	0.14	1.03	±	0.04